

Appendix 4. Characterization of site and treatment of DVT.

| | | Without PTS (n = 41) | With PTS (n = 60) | Total (n = 101) | p |
|--------------------------------------|------------|---------------------------------|------------------------------|----------------------------|----------|
| DVT Site | | | | | |
| Iliac* n (%) | | 10 (24) | 12 (20) | 22 (22) | NS |
| Femoral* | | 20 (49) | 34 (57) | 54 (54) | NS |
| Popliteal* | | 24 (59) | 41 (68) | 65 (64) | NS |
| Gemelar* | | 10 (24) | 10 (17) | 20 (20) | NS |
| Number of segments involved* | 1 | 23 (56) | 27 (45) | 50 (50) | NS |
| | 2 | 13 (32) | 26 (43) | 39 (39) | NS |
| | 3 | 5 (12) | 6 (10) | 11 (11) | NS |
| Side (right)* | | 15 (37) | 28 (47) | 43 (43) | NS |
| Pulmonary embolism | | 0 | 1 (2) | 1 (1) | NS |
| Elastic Compression Stockings | | | | | |
| Use of stocking | | 34 (83) | 50 (83) | 84 (83) | NS |
| Type of Stocking | Thigh-high | 27 (66) | 41 (68) | 68 (67) | NS |
| | Knee-high | 7 (17) | 9 (15) | 16 (16) | NS |
| Duration (months), median (IQR) | | 6 (11) | 5 (20) | 6 (12) | NS |
| Time per day (hours), median (IQR) | | 14 (5) | 12 (14) | 12 (14) | NS |
| Anticoagulation Therapy * | | | | | |
| Enoxoparin | | 41 (100) | 59 (98) | 100 (99) | NS |
| Warfarin | | 39 (95) | 58 (97) | 97 (96) | NS |
| Acenocumarol | | 2 (5) | 0 | 2 (2) | NS |
| Duration (months), median (IQR) | | 6 (7) | 12 (19) | 6 (13) | NS |

Legend: *Unknown in 1 patient with presence of PTS; IQR – Interquartile range; NS - Not significant.