

## Arrhythmogenic Left Ventricular Cardiomyopathy: A Successful Case of Extracorporeal Cardiopulmonary Resuscitation

### Miocardíopatia Arritmogénica do Ventrículo Esquerdo: Um Caso de Sucesso de Ressuscitação Cardiopulmonar Extracorporeal

Mafalda GAMA<sup>1</sup>, Isabel CARDOSO<sup>2</sup>, Mónica PALMA ANSELMO<sup>3</sup>, Sílvia AGUIAR ROSA<sup>2</sup>, Pedro GASPARD DA COSTA<sup>1</sup>, Philip FORTUNA<sup>1</sup>

Acta Med Port 2023 Sep;36(9):598-602 • <https://doi.org/10.20344/amp.19624>

#### ABSTRACT

A 24-year-old man suffered a witnessed cardiac arrest after a padel game. Basic life support was immediately provided. The pre-hospital emergency services team continued the resuscitation efforts, and the patient was accepted for extracorporeal cardiopulmonary resuscitation. The return of spontaneous circulation was achieved in 45 minutes. The initial assessment revealed a ST-segment elevation in leads V<sub>4</sub>-V<sub>6</sub> and a dilated left ventricle with severe systolic dysfunction. Coronary angiography was normal. An improvement in left ventricular systolic function was observed and extracorporeal cardiac support was discontinued after 48 hours. Cardiovascular magnetic resonance imaging demonstrated hypokinesia and subepicardial fatty infiltration of the left ventricle lateral wall. Genetic testing detected a variant of uncertain significance in the ANK2 gene. The diagnosis of arrhythmogenic left ventricular cardiomyopathy did not fulfill all the current diagnostic criteria, but it is a very likely diagnosis. An implantable cardioverter-defibrillator was placed. The patient was discharged without physical or cognitive impairment.

**Keywords:** Arrhythmogenic Left Ventricular Cardiomyopathy; Cardiopulmonary Resuscitation; Extracorporeal Membrane Oxygenation; Heart Arrest/therapy

#### RESUMO

Homem de 24 anos sofreu uma paragem cardiorrespiratória presenciada após um jogo de padel. O suporte básico de vida foi imediatamente iniciado. A equipa de emergência pré-hospitalar manteve os esforços de ressuscitação e o doente foi aceite para ressuscitação cardiopulmonar extracorporeal. O retorno da circulação espontânea foi atingido de imediato após 45 minutos. A avaliação inicial evidenciou elevação do segmento ST nas derivações V<sub>4</sub>-V<sub>6</sub> e dilatação do ventrículo esquerdo com disfunção sistólica grave. Angiografia coronária sem alterações. Foi observada uma melhoria da função sistólica do ventrículo esquerdo e a oxigenação por membrana extracorporeal veno-arterial foi suspensa após 48 horas. A ressonância magnética cardiovascular demonstrou hipocinesia da parede lateral e infiltração gordurosa subepicárdica na parede lateral do ventrículo esquerdo. O teste genético revelou uma variante de significado incerto no gene ANK2. Apesar do diagnóstico de miocardíopatia arritmogénica do ventrículo esquerdo não preencher todos os critérios diagnósticos atuais, é, no entanto, um diagnóstico muito provável. Foi colocado um cardioversor desfibrilador implantável. O doente teve alta sem compromisso físico ou cognitivo.

**Palavras-chave:** Miocardíopatia Arritmogénica do Ventrículo Esquerdo; Oxigenação por Membrana Extracorporeal; Paragem Cardíaca/tratamento Ressuscitação Cardiopulmonar

#### INTRODUCTION

Sudden cardiac death is a devastating event that occurs unexpectedly. A significant number of cases are attributed to non-coronary causes with a genetic basis, such as cardiomyopathies, channelopathies and malignant arrhythmias.

The arrhythmogenic cardiomyopathy (ACM) is a heart muscle disease that affects the right ventricle (RV), the left ventricle (LV) or both, according to the 2020 International criteria for ACM. It is characterized structurally by a myocardial scar (fibro or fibrofatty myocardial replacement) and functionally by ventricular dysfunction. This myocardial scar is a predisposing factor for ventricular arrhythmias, which could present as sudden cardiac arrest, regardless of the severity of pump failure.<sup>1,2</sup>

The diagnosis of ACM is made by meeting the Padua

criteria. A definitive diagnosis of arrhythmogenic left ventricular cardiomyopathy (ALVC) must meet one major structural LV criterion along with the presence of a gene mutation for ACM.<sup>1,3</sup>

A complete understanding of the genetic basis of certain cardiac conditions is an area of medical knowledge that has significant potential for development. Advances in genomic technology and research have allowed for the identification of specific genes and mutations that play a role in the development of these conditions.

The authors consider that this case report serves as an important warning about the association between mutations in the ANK2 gene with structural changes in the myocardium, which can increase the likelihood of malignant arrhythmias and sudden death.

1. Unidade de Urgência Médica. Centro Hospitalar Universitário de Lisboa Central. Lisbon. Portugal.

2. Serviço de Cardiologia. Centro Hospitalar Universitário de Lisboa Central. Lisbon. Portugal.

3. Hospital Professor Doutor Fernando da Fonseca. Lisbon. Portugal.

✉ Autor correspondente: Mafalda Gama. [mafalda.agm@gmail.com](mailto:mafalda.agm@gmail.com)

Recebido/Received: 30/01/2023 - Aceite/Accepted: 03/05/2023 - Publicado Online/Published Online: 16/06/2023 - Publicado/Published: 01/09/2023

Copyright © Ordem dos Médicos 2023



## CASE REPORT

We describe a case of a 24-year-old man, without relevant personal or family history. The individual suffered a witnessed cardiac arrest after a padel game. Basic life support (BLS) was immediately provided by laypeople. Eight minutes later the pre-hospital emergency services team arrived and continued the resuscitation efforts in accordance with advanced life support (ALS) best practice. The initial arrest rhythm was ventricular fibrillation (VF). The field team established early liaison with the Intensive Care Unit (ICU), and the patient was accepted for extracorporeal cardiopulmonary resuscitation (eCPR). During transport, quality mechanical chest compressions were assured. The patient arrived at the ICU 40 minutes after the onset of cardiac arrest in refractory VF. The right common femoral artery and vein were cannulated percutaneously with ultrasound guidance, and veno-arterial extracorporeal membrane oxygenation (VA-ECMO) was started within five minutes. The return of spontaneous circulation was achieved 45 minutes after the onset of cardiac arrest in sinus rhythm (with eight minutes of BLS, 32 minutes of ALS, and five minutes for cannulation to establish VA-ECMO). For distal limb perfusion, the ipsilateral superficial femoral artery was cannulated.

The initial 12-lead electrocardiogram presented with a ST-segment elevation in leads  $V_4$ - $V_6$  (Fig. 1). Transthoracic echocardiography (TTE) revealed a dilated LV with severe systolic dysfunction due to global diffuse hypokinesia. The cardiac output was severely depressed [left ventricular outflow tract velocity time integral (LVOT VTI) of 4.8 cm], a non-dilated RV with a mild systolic dysfunction on visual assessment. Coronary angiography revealed normal coronary

vessels.

For neuroprotection, the patient underwent deep sedation monitored by continuous electroencephalogram and therapeutic temperature management (33°C) with an external cooling device in the first 24 hours.

Initially, the patient was kept on VA-ECMO support (blood flow of 2.5 – 3.0 L/minute and sweep gas flow of 1 L/minute). Limb perfusion was monitored with bilateral near-infrared spectroscopy.

After 24 hours of neuroprotection bundle, the patient was slowly reheated to normothermia. With sedative reduction, the patient had spontaneous eye opening and was able to follow the observer and obey commands.

As cardiac function improved, VA-ECMO support was slowly reduced with no cardiovascular dysfunction or perfusion imbalance. TTE revealed an improvement of LVOT VTI to 17.6 cm and RV function (tricuspid annular plane systolic excursion of 18 mm and S' RV of 11 mm). A mild acute renal injury was observed in the acute phase and was rapidly resolved.

After 48 hours of VA-ECMO, the patient was successfully weaned and decannulated. Respiratory function was normal, and he was successfully extubated. No other organ support was required. Repeated clinical neurological examination revealed a progressive improvement in brain function to a cerebral performance category (CPC) score of 1 (CPC 1 – good cerebral performance).

Four days after the cardiac arrest, the patient was transferred to the Cardiology ward for further studies to clarify the cause of cardiac arrest.

Cardiovascular magnetic resonance imaging (CMR)




Figure 1 – Initial 12-lead electrocardiogram presented with a ST-segment elevation from  $V_4$  to  $V_6$




demonstrated non-dilated moderately impaired LV [LV ejection fraction (EF) of 48%], hypokinesia of the lateral wall and subepicardial fatty infiltration of the LV lateral wall. T2 mapping was increased in the lateral walls suggesting myocardial oedema. Extensive subepicardial late gadolinium enhancement (LGE) was noted in the anterior, lateral and inferior walls. Preserved systolic function of RV (RV EF 51%) with no regional wall motion abnormalities and no LGE in the RV wall was also noted (Fig. 2).

Based on the clinical history and the imaging findings, ALVC was the most likely diagnosis. Genetic tests were per-

formed. An implantable cardioverter-defibrillator (ICD) was placed for secondary prevention of sudden cardiac death.

The patient was discharged from the hospital on the 18<sup>th</sup> day after cardiac arrest. The patient demonstrated no physical or cognitive impairment and was found to be completely independent (with Medical Research Council sum-score of 60 – normal strength) and CPC 1.<sup>4</sup> The 12-lead electrocardiogram at the time of discharge showed no abnormalities (Fig. 3). The patient received follow-up in the cardiomyopathy clinic and in the following seven months there were no arrhythmic events.



**Figure 2** – Cardiovascular magnetic resonance imaging showing subepicardial fatty infiltration of the left ventricular lateral wall (arrows) in cine images (A) and native T1 mapping (B); extensive subepicardial late gadolinium enhancement in the anterior, lateral and inferior walls (C and D)




Figure 3 – The 12-lead electrocardiogram at discharge without alterations

Genetic testing detected a variant c.11081A>G, p.(Lys3694Arg), in the *ANK2* gene which is classified as a variant of uncertain significance. The diagnosis of ALVC does not fulfill all the current diagnostic criteria.

## DISCUSSION

This case represents a successful case of eCPR in refractory cardiac arrest. The following factors predicted a high probability of survival with favorable neurological outcome: younger age, no comorbidities, witnessed arrest, no-flow interval of less than five minutes, initial cardiac rhythm of VF, transportation with high quality chest compressions, refractory VF and low-flow interval of less than 60 minutes. For the same reason, those are typically considered the inclusion criteria for admission to eCPR.<sup>5</sup>

This case is compatible with the diagnosis of ALVC due to the clinical picture of recovered cardiac arrest after a refractory VF, imaging findings with mild LV systolic dysfunction and myocardial scar on CMR, and exclusion of ischemic, valvular, or infectious disease. Genetic testing did not fully confirm the hypothesis since it is a poorly described mutation without definitive clinical relevance.

The *ANK2* gene mutation was initially identified as a causative mutation for long QT syndrome and later for Brugada syndrome.<sup>6,7</sup> Interestingly, the *ANK2* gene mutations have also been found in individuals with structural myocardial changes, even without a prolonged QT interval, which increases their susceptibility to malignant arrhythmias and sudden death.<sup>8,9</sup> Therefore, additional genotyping and phenotyping studies are necessary and encouraged.

According to the International criteria for ACM, it is mandatory to have ACM-related genetic mutations associated with LV structural abnormalities to confirm the diagnosis.<sup>1</sup> In fact, morpho-functional and structural LV abnormalities of ACM do not provide sufficient disease specificity because of the overlap with the phenotypic features of other heart muscle diseases, and hence the stricter criteria.<sup>3</sup>

This patient fulfilled one major criteria (LV LGE) and two minor criteria (global LV systolic dysfunction and regional LV hypokinesia of LV free wall). Therefore, the diagnosis of ALVC was not confirmed, although it is very likely.<sup>2</sup>

As the diagnostic criteria for arrhythmogenic cardiomyopathies have changed with the development of knowledge about the diseases, the diagnostic criteria that define ALVC may change, or other genetic mutations related to ACM may be found in the future.

These types of conditions do not have a specific treatment and ICD implantation is the only strategy that may prevent sudden cardiac death.

The authors consider this case report helpful to other clinicians or researchers in various aspects. First, we demonstrated a successful case of an eCPR in the presence of adequate patient selection, teamwork and appropriate interventions in the different stages of patient care. Second, it adds to the literature by highlighting a possible additional genetic pathway to be explored and researched in future cases of cardiac arrest, arrhythmia, heart muscle diseases and ALVC.

**AUTHOR CONTRIBUTIONS**

MG: Conceptualization, writing, review and editing of the final version of the manuscript.

IC, MPA, SAR, PGC, PF: Writing, review and editing of the manuscript.

**PROTECTION OF HUMANS AND ANIMALS**

The authors declare that the procedures were followed according to the regulations established by the Clinical Research and Ethics Committee and to the Helsinki Declaration of the World Medical Association updated in 2013.

**DATA CONFIDENTIALITY**

The authors declare having followed the protocols in use at their working center regarding patients' data publication.

**REFERENCES**

1. Corrado D, Zorzi A, Cipriani A, Bauce B, Bariani R, Boffagna G, et al. Evolving diagnostic criteria for arrhythmogenic cardiomyopathy. *J Am Heart Assoc.* 2021;10:e021987.
2. Corrado D, Perazzolo Marra M, Zorzi A, Boffagna G, Cipriani A, Lazzari M, et al. Diagnosis of arrhythmogenic cardiomyopathy: the Padua criteria. *Int J Cardiol.* 2020;319:106-14.
3. Corrado D, Basso C. Arrhythmogenic left ventricular cardiomyopathy. *Heart.* 2022;108:733-43.
4. Hermans G, Clerckx B, Vanhullebusch T, Segers J, Vanpee G, Robbeets C, et al. Interobserver agreement of Medical Research Council sum-score and handgrip strength in the intensive care unit. *Muscle Nerve.* 2012;45:18-25.
5. Richardson AS, Tonna JE, Nanjaya V, Nixon P, Abrams DC, Raman L, et al. Extracorporeal cardiopulmonary resuscitation in adults. Interim guideline consensus statement from the Extracorporeal Life Support Organization. *ASAIO J.* 2021;67:221-8.
6. Mohler PJ, Schott JJ, Gramolini AO, Dilly KW, Guatimosim S, duBell WH, et al. Ankyrin-B mutation causes type 4 long-QT cardiac arrhythmia and sudden cardiac death. *Nature.* 2003;421:634-9.
7. Allegue Toscano C, Coll Vidal M, Matés Ramírez J, Campuzano Larrea O, Iglesias A, Sobrino B, et al. Genetic analysis of arrhythmogenic diseases in the era of NGS: the complexity of clinical decision-making in brugada syndrome. *PLoS One.* 2015;7:e0133037.
8. Roberts JD, Murphy NP, Hamilton RM, Lubbers ER, James CA, Kline C, et al. Ankyrin-B dysfunction predisposes to arrhythmogenic cardiomyopathy and is amenable to therapy. *J Clin Investig.* 2019;129:3171-84.
9. Mohler PJ, Splawski I, Napolitano C, Bottelli G, Sharpe L, Timothy K, et al. A cardiac arrhythmia syndrome caused by loss of ankyrin-B function. *Proc Natl Acad Sci.* 2004;101:9137-42.

**PATIENT CONSENT**

Obtained.

**COMPETING INTERESTS**

PF has received support for attending meetings and/or travel from Hamilton Medical and Getinge.

All other authors have declared that no competing interests exist.

**FUNDING SOURCES**

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.